





	7011		U	AL AM	len	(of Medicine NI	M.	
PubMed	Nucleo	otide	Protein	Genome	Structure	PMC	Taxonomy	ОМІМ	Во
Search Publ	1ed	Y	for				Go Clear		
			Limits	Preview/Inde	x Histo	ory	Clipboard	Det	ails
About Entrez	J	Dist	olay Abstr	act 🎼	Show: 20	Sor	t 🕞 Ser	id to Te	ext
Text Version			,		,				************
		□1:	Ann Thora	c Surg. 1998 M	ay;65(5):131	3-5.		Related A	Articles

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Comment in:

Ann Thorac Surg. 1998 Dec;66(6):2158-9.

PENEVIER CHENCE
FULL-TEXT ARTICLE

Aortic valve operations under deep hypothermic circulatory arr for the porcelain aorta: "no-touch" technique.

Byrne JG, Aranki SF, Cohn LH.

Division of Cardiac Surgery, Brigham and Women's Hospital, Harvard Medi School, Boston, Massachusetts 02115, USA.

BACKGROUND: Aortic valve replacement or repair becomes a high-risk procedure in patients in whom the ascending aorta cannot be clamped either because of extensive calcification and risk of cerebral embolus or because of extensive adhesions precluding safe dissection and clamping. METHODS: W report the results of aortic valve replacement or repair with deep hypothermic circulatory arrest in 3 patients. Techniques to improve results include routine of epiaortic and transesophageal echocardiography, avoidance of manipulatio the ascending aorta until the circulation is arrested, avoidance of antegrade cardioplegia, routine use of retrograde cardioplegia and retrograde cerebral perfusion, when feasible, and minimal aortotomy (just enough to excise and replace or repair the valve). RESULTS: Operations were accomplished in approximately 1 hour each with minimal manipulation of the aorta, thus minimizing aortic trauma and subsequent risk of cerebral embolus. Each patie had an unremarkable recovery without neurologic complications. CONCLUSIONS: Aortic valve replacement or repair using the "no-touch" technique and deep hypothermic circulatory arrest is the preferred method wh dealing with the porcelain or unclampable aorta.

PMID: 9594858 [PubMed - indexed for MEDLINE]

Display Abstract 💌	Show: 20 Sort	Send to Text

Write to the Help Desk

NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Jul 17 2003